

# ON CALCULUS WITH A QUATERNIONIC VARIABLE AND ITS CHARACTERISTIC CAUCHY-RIEMANN TYPE EQUATIONS

DANIEL ALAYÓN-SOLARZ

ABSTRACT. We show sufficient and necessary conditions, in terms of partial differential equations with variable coefficients, for a quaternionic function to admit a continuous derivative in a open set in the sense of C. Schwartz.

## 1. INTRODUCTION

Let  $\mathbb{H}$  denote the quaternions, let  $q = t + xi + yj + zk$  be a quaternion written as  $q = t + r\iota$ , where  $t$  is the real part of the quaternion,  $r \geq 0$  and  $\iota^2 = -1$ . The problem of the quaternionic left-derivative starts by considering the limit

$$\lim_{h \rightarrow 0} h^{-1}(f(q+h) - f(q) - hA) = 0$$

for some number  $A$ . It is well known that the only functions left-derivable in  $\mathbb{H}$  in the above sense turn out to be  $f(q) = qa + b$  for some  $a, b \in \mathbb{H}$ . A new approach for this problem, which results in a larger class of functions has been introduced in [2].

**Definition 1.** Let  $f$  be a quaternionic function that is continuous and real differentiable at  $q$ . We say  $f$  is  $S$ -derivable by the left at  $q$  if there exists quaternions  $A$  or  $B$  and  $C$  such that the limit

$$\lim_{h \rightarrow 0} \begin{cases} h^{-1}(f(q+h) - f(q) - hA) & \text{if } q \in \mathbb{R} \\ h^{-1}(f(q+h) - f(q) - h_{\parallel}B - h_{\perp}C) & \text{if } q \in \mathbb{H} \setminus \mathbb{R} \end{cases}$$

where

$$h_{\parallel} := \frac{h - \iota h \iota}{2}, \quad h_{\perp} := \frac{h + \iota h \iota}{2}$$

vanishes at  $q$ . Then we say the function  $f$  is  $S$ -derivable at  $q$ . The numbers  $A$  and  $B$  we shall call the parallel derivative and  $C$  the perpendicular derivative at  $q$ .

Just like in the complex setting we would like to consider functions that are derivable in this new, broader sense in some open set  $\Omega \subseteq \mathbb{H}$ .

**Definition 2.** Let  $f : \Omega \subseteq \mathbb{H} \rightarrow \mathbb{H}$  be a quaternionic function that is continuously real-differentiable. We say  $f$  is continuously  $S$ -derivable by the left in  $\Omega$  if there exists continuous quaternionic functions  $A(q) : \Omega \cap \mathbb{R} \rightarrow \mathbb{H}$  and  $B(q), C(q) : \Omega \setminus \mathbb{R} \rightarrow \mathbb{H}$  such that

$$\lim_{h \rightarrow 0} \begin{cases} h^{-1}(f(q+h) - f(q) - hA(q)) = 0 & \text{if } q \in \mathbb{R} \\ h^{-1}(f(q+h) - f(q) - h_{\parallel}B(q) - h_{\perp}C(q)) = 0 & \text{if } q \in \mathbb{H} \setminus \mathbb{R} \end{cases}$$

for all  $q \in \Omega \subseteq \mathbb{H}$ .

In the complex case one obtains characteristic equations, namely the Cauchy-Riemann equations by evaluation of the derivative with two prescribed directions, the real and the pure imaginary. The result we consider first that this class can be characterized by *some* partial differential equations, just like in the commutative complex setting. More precisely:

**Theorem 1.** *A function quaternionic function  $f$  whose partial derivatives are continuous in some open set  $\Omega$ , then  $f$  is continuously  $S$ -derivable in  $\Omega$  if, and only if, it satisfies the following partial differential equations in  $\Omega \cap \mathbb{R}$*

$$\frac{\partial f}{\partial x} = i \frac{\partial f}{\partial t} \quad (1)$$

$$\frac{\partial f}{\partial y} = j \frac{\partial f}{\partial t} \quad (2)$$

$$\frac{\partial f}{\partial y} = k \frac{\partial f}{\partial t} \quad (3)$$

and the following partial differential equations in  $\Omega \setminus \mathbb{R}$

$$\frac{\partial f}{\partial x} = \left( \frac{i - i\iota}{2} \right) \frac{\partial f}{\partial t} - \left( \frac{i + i\iota}{4} \right) Df \quad (4)$$

$$\frac{\partial f}{\partial y} = \left( \frac{j - j\iota}{2} \right) \frac{\partial f}{\partial t} - \left( \frac{j + j\iota}{4} \right) Df \quad (5)$$

$$\frac{\partial f}{\partial z} = \left( \frac{k - k\iota}{2} \right) \frac{\partial f}{\partial t} - \left( \frac{k + k\iota}{4} \right) Df \quad (6)$$

where  $D$  denotes the left Fueter operator:

$$Df := \frac{\partial f}{\partial t} + i \frac{\partial f}{\partial x} + j \frac{\partial f}{\partial y} + k \frac{\partial f}{\partial z}$$

This equations clearly play here the role of the Cauchy-Riemann equations played in the commutative setting. We then show that if a function is  $S$ -derivable then is necessarily Cullen-regular (also termed Slide regular) in the sense of G. Gentili and D.C Struppa. Cullen-regular functions have been shown, see [1] to be expressible as absolutely convergent quaternionic unilateral power series assuming the domain of the function is a ball centered at the origin with radius  $R$ . One consequence of the previous result is that it allows to state the following characterization of a continuously  $S$ -derivable function in a ball of radius  $R$  centered at the origin.

**Theorem 2.** *Let  $f : B(0, R) \rightarrow \mathbb{H}$  be a  $C^1$  quaternionic function. Then  $f$  is  $S$ -derivable by the left if and only if it is expressible as a absolutely convergent power series*

$$f(q) = \sum_{k=0}^{\infty} q^k a_k, \quad a_k \in \mathbb{H}$$

and

$$A(q) = \frac{\partial f}{\partial t}, \quad q \in B(0, R) \cap \mathbb{R}$$

$$B(q) = \frac{\partial f}{\partial t}, \quad q \in B(0, R) \cap \mathbb{H} \setminus \mathbb{R}$$

$$C(q) = (q - \bar{q})^{-1}[f(q) - f(\bar{q})]$$

Thus recovering the original result of C. Schwartz.

## 2. NECESSARY CONDITION FOR S-DIFFERENTIABILITY USING DIRECTIONS BASED ON $1, i, j$ AND $k$

In complex calculus the necessity of the Cauchy-Riemann equations is usually obtained by considering the limit that defines the derivative in the real and pure imaginary directions at a point. Proceeding analogously with the quaternionic case one obtains corresponding identities.

**Proposition 1.** *Let  $f$  be continuously S-differentiable in  $\Omega$ , then*

$$A(q) = \frac{\partial f}{\partial t}(q), q \in \Omega$$

*Proof.* By the nature of the definition we must check the cases when the base point  $q$  is real or not. Let  $q$  be real and  $f$  S-derivable at  $q$ . Choosing  $h$  as a real number and after taking the limit as  $h \rightarrow 0$  we obtain

$$A(q) = \frac{\partial f}{\partial t}(q)$$

Now assume  $q$  is not real. Choose again  $h$  a real number. Since  $h$  is real then  $h_{\parallel} = h$  and  $h_{\perp} = 0$ , so if  $f$  is S-differentiable we have

$$B(q) = \frac{\partial f}{\partial t}(q)$$

for all  $q \in \Omega \setminus \mathbb{R}$ . □

**Proposition 2.** *Let  $f$  be continuously S-differentiable, then  $f$  satisfies equations (1) to (6) and*

$$C(q) = -\left(\frac{1}{2}\right)Df(q)$$

where  $D$  is the left Fueter operator.

*Proof.* The preceding proposition means  $A(q)$  and  $B(q)$  can be replaced by  $\frac{\partial f}{\partial t}$  without loss of generality. Again we start with  $q$  real. Choosing  $h$  as  $\epsilon i$ ,  $\epsilon j$  and  $\epsilon k$  where  $\epsilon$  is real and taking the corresponding limit as  $\epsilon$  goes to zero we obtain necessary conditions for the partial derivatives with respect to  $x, y$  and  $z$ . If  $q$  is real we obtain immediately equations (1), (2) and (3). If  $q$  is non-real then we obtain the following equations, with  $C(q)$  a function to be determined:

$$\frac{\partial f}{\partial x} = \left(\frac{i - \iota i}{2}\right) \frac{\partial f}{\partial t} + \left(\frac{i + \iota i}{2}\right) C(q), \quad (7)$$

$$\frac{\partial f}{\partial y} = \left(\frac{j - \iota j}{2}\right) \frac{\partial f}{\partial t} + \left(\frac{j + \iota j}{2}\right) C(q), \quad (8)$$

$$\frac{\partial f}{\partial z} = \left(\frac{k - \iota k}{2}\right) \frac{\partial f}{\partial t} + \left(\frac{k + \iota k}{2}\right) C(q), \quad (9)$$

For this we multiply Eq. (7) by the left by  $i$ , Eq. (8) by the left by  $j$  and Eq. (9)

by the left by  $k$  and sum all three resulting equations. For the left hand side one thus obtains:

$$i \frac{\partial f}{\partial x} + j \frac{\partial f}{\partial y} + k \frac{\partial f}{\partial z}.$$

The right hand side is further simplified with the following identities:

$$i \left( \frac{i - \iota \iota}{2} \right) + j \left( \frac{j - \iota j \iota}{2} \right) + k \left( \frac{k - \iota k \iota}{2} \right) = -1,$$

and

$$i \left( \frac{i + \iota \iota}{2} \right) + j \left( \frac{j + \iota j \iota}{2} \right) + k \left( \frac{k + \iota k \iota}{2} \right) = -2.$$

Thus one obtains

$$i \frac{\partial f}{\partial x} + j \frac{\partial f}{\partial y} + k \frac{\partial f}{\partial z} = -\frac{\partial f}{\partial t} - 2C(q)$$

or

$$\frac{\partial f}{\partial t} + i \frac{\partial f}{\partial x} + j \frac{\partial f}{\partial y} + k \frac{\partial f}{\partial z} = -2C(q)$$

□

### 3. SUFFICIENT CONDITIONS FOR THE S-DERIVABILITY

**Proposition 3.** *Let  $f : \Omega \rightarrow \mathbb{H}$  be continuously real-differentiable and suppose that  $f$  satisfies Eq. (1) to (6) in  $\Omega$ , then  $f$  is S-derivable in  $\Omega$ .*

*Proof.* Because  $f$  is continuously real differentiable in  $\Omega$  then:

$$\lim_{h \rightarrow 0} \frac{|f(q+h) - f(q) - M_f h|}{|h|} = 0$$

where  $M_f$  is the Jacobi matrix of  $f$  at  $q$ . Now writing  $h = h_0 + ih_1 + jh_2 + kh_3$ ,  $h_n \in \mathbb{R}$  we can also write

$$M_f h = \frac{\partial f}{\partial t} h_0 + \frac{\partial f}{\partial y} h_1 + \frac{\partial f}{\partial y} h_2 + \frac{\partial f}{\partial z} h_3$$

If  $q$  is real one starts with and use Eq. (1), (2) and (3), to obtain

$$M_f h = h \frac{\partial f}{\partial t}.$$

For  $q$  non real, and if  $f$  satisfies equations (4),(5) and (6) we have

$$\begin{aligned} & \frac{\partial f}{\partial t} h_0 + \frac{\partial f}{\partial y} h_1 + \frac{\partial f}{\partial y} h_2 + \frac{\partial f}{\partial z} h_3 = \\ & \frac{\partial f}{\partial t} h_0 + \left( \frac{ih_1 - \iota h_1 \iota}{2} \right) \frac{\partial f}{\partial t} - \left( \frac{ih_1 + \iota h_1 \iota}{4} \right) Df \\ & + \left( \frac{jh_2 - \iota jh_2 \iota}{2} \right) \frac{\partial f}{\partial t} - \left( \frac{jh_2 + \iota jh_2 \iota}{4} \right) Df \\ & + \left( \frac{kh_3 - \iota kh_3 \iota}{2} \right) \frac{\partial f}{\partial t} - \left( \frac{kh_3 + \iota kh_3 \iota}{4} \right) Df \end{aligned}$$

which can be rewritten in the compact form

$$M_f h = h_{\parallel} \frac{\partial f}{\partial t} + h_{\perp} \left( \frac{-1}{2} Df \right)$$

therefore we conclude that for  $q$  real:

$$\lim_{h \rightarrow 0} |h^{-1} (f(q+h) - f(q) - h \frac{\partial f}{\partial t})| = \lim_{h \rightarrow 0} \frac{|f(q+h) - f(q) - M_f h|}{|h|} = 0$$

which implies

$$\lim_{h \rightarrow 0} h^{-1}(f(q+h) - f(q) - h \frac{\partial f}{\partial t}) = 0$$

and similarly, for a  $q$  non-real:

$$\lim_{h \rightarrow 0} |h^{-1}(f(q+h) - f(q) - h_{\parallel} \frac{\partial f}{\partial t} + h_{\perp} \frac{1}{2} Df)| = \lim_{h \rightarrow 0} \frac{|h(f(q+h) - f(q) - M_f h)|}{|h|} = 0$$

which implies

$$\lim_{h \rightarrow 0} h^{-1}(f(q+h) - f(q) - h_{\parallel} \frac{\partial f}{\partial t} + h_{\perp} \frac{1}{2} Df) = 0$$

and therefore we have found the  $A(q)$ ,  $B(q)$  and  $C(q)$  required for this function  $f$  to be continuously S-derivable.  $\square$

#### 4. CULLEN-REGULARITY AND COMMUTING DIRECTIONAL DERIVATIVES

The above section shows that the four directions  $1, i, j$  and  $k$  are sufficient and necessary for finding characteristic equations. However  $i, j$  and  $k$  are mere examples of  $\sqrt{-1}$  in  $\mathbb{H}$ . Let  $\mathbb{S}^2 \subseteq \mathbb{H}$  be the set of roots of  $-1$ , and fix  $I \in \mathbb{S}^2$ . Since  $I^2 = -1$  the set  $\mathbb{R} + I\mathbb{R}$  is a isomorphic copy of  $\mathbb{C}$ . Now let  $f$  be continuously S-derivable in some  $\Omega$  whose intersection with the real numbers is non-empty. For a real  $q$  we consider

$$\lim_{I\epsilon \rightarrow 0} (I\epsilon)^{-1}(f(q + \epsilon I) - f(q)) = \frac{\partial f}{\partial t}$$

on the other side, interpreting the left-hand side as a directional derivative:

$$\lim_{\epsilon \rightarrow 0} \frac{f(q + \epsilon I) - f(q)}{\epsilon} = \frac{\partial f}{\partial r}$$

when restricted to  $\mathbb{R} + I\mathbb{R}$ . Thus we conclude that for  $q$  a real number the expression

$$\frac{\partial f}{\partial t} + I \frac{\partial f}{\partial r} = 0$$

vanishes in  $\mathbb{R} + I\mathbb{R}$ . If  $q \in \mathbb{H} \setminus \mathbb{R}$ , write it as  $q = t + rI$ , for some  $I^2 = -1$  and  $r \geq 0$ . Consider increments only in the imaginary component of  $\mathbb{R} + I\mathbb{R}$ :  $h = \epsilon I$ , then  $h_{\parallel} = \epsilon I$ ,  $h_{\perp} = 0$ . Then

$$\frac{\partial f}{\partial r} = \lim_{\epsilon \rightarrow 0} \frac{f(q + \epsilon I) - f(q)}{\epsilon} = I \frac{\partial f}{\partial t}$$

and, as  $I^2 = -1$ :

$$\frac{\partial f}{\partial t} + I \frac{\partial f}{\partial r} = 0$$

at  $q$ . Let  $\iota : \mathbb{H} \setminus \mathbb{R} \rightarrow \mathbb{S}^2 \subseteq \mathbb{H} \setminus \mathbb{R}$  be the function that sends a quaternion  $q = t + rI$  to  $I$ . Then the above means that  $f$  satisfies

$$(\frac{\partial}{\partial t} + \iota \frac{\partial}{\partial r})f = 0.$$

This discussion shows that S-derivable functions are necessarily Cullen-regular in the sense of G. Gentili and D.C Struppa.

## 5. THE IOTA-DERIVATIVE AND THE PERPENDICULAR DERIVATIVE

The previous section only considered increments that commuted with the variable  $q$ . Start parametrizing the function  $\iota$  as  $\cos \alpha \sin \beta i + \sin \alpha \sin \beta j + \cos \beta$  writing the left Fueter operator as

$$Df = \frac{\partial f}{\partial t} + \iota \frac{\partial f}{\partial r} - \frac{1}{r} \frac{\partial f}{\partial \iota}$$

where

$$\frac{\partial f}{\partial \iota} := \left(\frac{\partial \iota}{\partial \alpha}\right)^{-1} \frac{\partial}{\partial \alpha} + \left(\frac{\partial \iota}{\partial \beta}\right)^{-1} \frac{\partial}{\partial \beta}$$

Which is defined outside the singular subplane  $\mathbb{R} + k\mathbb{R}$ . observe that  $(\frac{\partial \iota}{\partial \alpha})^{-1}$  and  $(\frac{\partial \iota}{\partial \beta})^{-1}$  anti-commute with the function  $\iota$ , so they are geometrically perpendicular to  $\iota$ . For any continuously real-derivable Cullen-regular function  $f$  we can write

$$Df = -\frac{1}{r} \frac{\partial f}{\partial \iota}$$

so the perpendicular derivative can also be written, without loss of generality as

$$\frac{1}{2r} \frac{\partial f}{\partial \iota}.$$

## 6. A CRITERIUM FOR THE S-DERIVABILITY

**Theorem 3.** *Let  $f : \Omega \rightarrow \mathbb{H}$  be a real-derivable quaternionic function such that*

- $f$  is  $S$ -derivable in  $\Omega \cap \mathbb{R}$ .
- $f = u + \iota v$  for some  $C^1$  quaternionic functions  $u, v$ .
- $u$  and  $v$  are independent of  $\alpha$  and  $\beta$ .
- $u$  and  $v$  satisfy:

$$\frac{\partial u}{\partial t} - \frac{\partial v}{\partial r} = \frac{\partial u}{\partial r} + \frac{\partial v}{\partial t} = 0$$

$$\Omega \setminus \mathbb{R}$$

then with this conditions  $f$  is continuously left  $S$ -derivable in  $\Omega$  and its perpendicular derivative is  $\frac{v}{r}$ .

*Proof.* Let  $f$  satisfy the hypothesis. We show  $f$  is continuously  $S$ -derivable in  $\Omega \setminus \mathbb{R}$ . Computing the partial derivatives:

$$\begin{aligned} \frac{\partial f}{\partial x} &= \frac{\partial u}{\partial x} + \iota_x v + \iota \frac{\partial v}{\partial x} \\ \frac{\partial f}{\partial y} &= \frac{\partial u}{\partial y} + \iota_y v + \iota \frac{\partial v}{\partial y} \\ \frac{\partial f}{\partial z} &= \frac{\partial u}{\partial z} + \iota_z v + \iota \frac{\partial v}{\partial z} \end{aligned}$$

on the other hand, since neither  $u$  nor  $v$  depend on  $\alpha$  and  $\beta$  implies that

$$\begin{aligned} \frac{\partial u}{\partial x} &= \frac{\partial u}{\partial r} \frac{x}{r}, \quad \frac{\partial v}{\partial x} = \frac{\partial v}{\partial r} \frac{x}{r} \\ \frac{\partial u}{\partial y} &= \frac{\partial u}{\partial r} \frac{y}{r}, \quad \frac{\partial v}{\partial y} = \frac{\partial v}{\partial r} \frac{y}{r} \\ \frac{\partial u}{\partial z} &= \frac{\partial u}{\partial r} \frac{z}{r}, \quad \frac{\partial v}{\partial z} = \frac{\partial v}{\partial r} \frac{z}{r} \end{aligned}$$

thus we can write

$$\begin{aligned}\frac{\partial f}{\partial x} &= \frac{x}{r} \frac{\partial f}{\partial r} + \iota_x v \\ \frac{\partial f}{\partial y} &= \frac{y}{r} \frac{\partial f}{\partial r} + \iota_y v \\ \frac{\partial f}{\partial z} &= \frac{z}{r} \frac{\partial f}{\partial r} + \iota_z v\end{aligned}$$

the hypothesis on  $f$  imply that we can replace  $\frac{\partial f}{\partial r}$  with  $\iota \frac{\partial f}{\partial t}$ .

$$\begin{aligned}\frac{\partial f}{\partial x} &= \frac{x}{r} \iota \frac{\partial f}{\partial t} + \iota_x v \\ \frac{\partial f}{\partial y} &= \frac{y}{r} \iota \frac{\partial f}{\partial t} + \iota_y v \\ \frac{\partial f}{\partial z} &= \frac{z}{r} \iota \frac{\partial f}{\partial t} + \iota_z v\end{aligned}$$

finally we need the following identities:

$$\frac{x}{r} \iota = \left( \frac{i - \iota i \iota}{2} \right), \quad \frac{y}{r} \iota = \left( \frac{j - \iota j \iota}{2} \right), \quad \frac{z}{r} \iota = \left( \frac{k - \iota k \iota}{2} \right)$$

and

$$\iota_x = \frac{1}{r} \left( \frac{i + \iota i \iota}{2} \right), \quad \iota_y = \frac{1}{r} \left( \frac{j + \iota j \iota}{2} \right), \quad \iota_z = \frac{1}{r} \left( \frac{k + \iota k \iota}{2} \right)$$

Since the function is Cullen-regular and neither  $u$  nor  $v$  depend of  $\alpha$  and  $\beta$  we conclude that

$$Df = -2 \frac{v}{r},$$

so  $f$  satisfies equations (4),(5) and (6).  $\square$

**Corollary 1.** *The function  $\iota$  is continuously S-derivable in  $\mathbb{H} \setminus \mathbb{R}$ .*

*Proof.* Take  $u = 0$  and  $v = 1$ .  $\square$

**Corollary 2.** *The power function is S-derivable in  $\mathbb{H}$ .*

*Proof.* Let  $f(q) = q^n$  for some  $n \in \mathbb{N}$ ,  $f$  is continuously real-derivable in  $\mathbb{H}$ . Let  $q$  be a real number. Then

$$\begin{aligned}\frac{\partial}{\partial t} q^n &= n q^{n-1} \\ \frac{\partial}{\partial x} q^n &= i n q^{n-1} \\ \frac{\partial}{\partial y} q^n &= j n q^{n-1} \\ \frac{\partial}{\partial z} q^n &= k n q^{n-1}\end{aligned}$$

and therefore  $q^n$  is S-derivable in  $\mathbb{R}$ . Since  $q^n = u_n(t, r) + \iota v_n(t, r)$ , for some real functions  $u_n$  and  $v_n$  satisfying

$$\frac{\partial u_n}{\partial t} - \frac{\partial v_n}{\partial r} = \frac{\partial u_n}{\partial r} + \frac{\partial v_n}{\partial t} = 0$$

we obtain the desired result.  $\square$

**Corollary 3.** *Let*

$$f(q) = \sum_{n=0}^{\infty} q^n a_n$$

*be an absolutely and uniformly convergent series in a ball  $B(0, R)$ . Then  $f$  is S-derivable in  $B(0, R)$ .*

*Proof.* Observe such  $f$  is real-derivable in  $B(0, 1)$ , (in fact is real-analytic). Let  $q \in B(0, R)$  be real. Uniform and absolute convergence means the series can be partial-derivated termwise. Thus one obtains that  $f$  satisfies equations (1),(2) and (3) for every  $q$  in  $B(0, R) \cap \mathbb{R}$ . On the other side we have

$$\sum_{n=0}^{\infty} q^n a_n = \sum_{n=0}^{\infty} u_n a_n + \iota \sum_{n=0}^{\infty} v_n a_n = u(t, r) + \iota v(t, r)$$

and such  $u$  and  $v$  satisfy:

$$\frac{\partial u}{\partial t} - \frac{\partial v}{\partial r} = \frac{\partial u}{\partial r} + \frac{\partial v}{\partial t} = 0$$

□

Observe that for functions constructed as in this section one has  $f(\bar{q}) = u(q) - \iota v(q)$ . So they also must satisfy the following equalities:

$$\frac{1}{2r} \frac{\partial f}{\partial \iota} = \frac{v}{r} = (\iota r)^{-1} (\iota v) = (q - \bar{q})^{-1} (f(q) - f(\bar{q}))$$

We are now ready to prove **Theorem 2**.

*Proof.* In [1] it is proved that every Cullen-regular functions  $f : B(0, R) \rightarrow \mathbb{H}$  can be expressed as such power series. Since S-derivable functions are in particular Cullen-regular this theorem remains valid. On the other side, functions expressible as such power series have been shown to be S-derivable themselves. The identities  $A(q) = B(q) = \frac{\partial f}{\partial \iota}$  and  $Df = -2(q - \bar{q})^{-1}(f(q) - f(\bar{q}))$  holds for all functions constructed as in **Theorem 3** and such power series are of this form. □

#### REFERENCES

- [1] G. Gentili, D.C. Struppa, *A new theory of regular functions of a quaternionic variable*, Adv. Math. **216**, 279-301. (2007)
- [2] C. Schwartz, *Calculus with a quaternionic variable* J. Math. Phys. **50**, 013523 (2009)

DEPARTAMENTO DE MATEMÁTICAS PURAS Y APLICADAS, UNIVERSIDAD SIMÓN BOLÍVAR, CARACAS, VENEZUELA.

*E-mail address:* danieldaniel@gmail.com